

Expansions of ${}_4F_3$ When the Upper Parameters Differ by Integers

MEGUMI SAIGO

Department of Applied Mathematics, Fukuoka University, Fukuoka 814-0180, Japan

R.K. SAXENA

Department of Mathematics and Statistics, Jai Narain Vyas University, Jodhpur-342001, India

(1991 AMS Classification number: 33C20)

In this article three expansion formulas for a generalized hypergeometric function ${}_4F_3$ are derived, when its upper parameters differ by integers. Though the results are special cases of a general continuation formula for ${}_pF_q$, they are sufficiently general and unify a number of known results.

1. Introduction

In many problems arising in physical sciences and statistics, hypergeometric functions, in logarithmic cases, are applicable, which is known from the monograph of Mathai and Haubold [4] and Mathai [3], etc. A detailed account of such applications is available from the two monographs by Mathai and Saxena [5], [6]. Expansions of the Gauss's hypergeometric functions in logarithmic cases are given in the monograph of Erdélyi *et al.* [1].

In a series of papers by Saigo *et al.* [9] - [25], numerous properties have been investigated which exhibits the behaviour of generalized hypergeometric functions near the boundaries of their regions of convergence of series defining these functions. In deriving some of these results, the expansions of Gauss's hypergeometric function in logarithmic cases given in the monograph by Magnus *et al.* [2] are used.

The object of this article is to derive three expansions of a generalized hypergeometric function ${}_4F_3(\cdot)$, when the upper parameters differ by integers. Though the results established in this article are special cases of a general continuation formula for ${}_pF_q$ [7, Entry 78 in §7.2.3], yet they are sufficiently general in nature and unify a number of known results in literature [1, p.63], [27, pp.41-43], etc. Further the results are obtained in a neat and compact form which may be used in investigating the asymptotic behaviour of the generalized hypergeometric functions.

From [8, p.101], we know that the generalized hypergeometric function ${}_4F_3(\cdot)$ has the following Mellin-Barnes integral representation

$$\begin{aligned} & \Gamma \left[\begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} \right] {}_4F_3 \left(\begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} ; z \right) \\ &= \frac{1}{2\pi i} \int_{\mathfrak{C}} \Gamma \left[\begin{matrix} a_1 + \xi, a_2 + \xi, a_3 + \xi, a_4 + \xi, -\xi \\ b_1 + \xi, b_2 + \xi, b_3 + \xi \end{matrix} \right] (-z)^\xi d\xi, \end{aligned} \quad (1.1)$$

where the contour $\mathfrak{C}$ is of Barnes type, extending from $-i\infty$ to $+i\infty$ in the ξ -plane, curving if necessary, such that the poles of gamma functions $\Gamma(a_1 + \xi)$, $\Gamma(a_2 + \xi)$, $\Gamma(a_3 + \xi)$ and $\Gamma(a_4 + \xi)$ lie to the left of the contour and the poles of

$\Gamma(-\xi)$ to its right.

In what follows m, n and p are non-negative integers and the notation

$$\Gamma \left[\begin{matrix} a_1, a_2, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} \right] = \frac{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)\Gamma(a_4)}{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)} \quad (1.2)$$

is employed.

The proofs of the results in §2 - §4 can be developed by an appeal to the calculus of residues and following a procedure described in [5].

2. First Expansion

We establish first the formula when two of the upper parameters differ by integers:

$$\begin{aligned} & \Gamma \left[\begin{matrix} a_1, a_1 + m, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} \right] {}_4F_3 \left(\begin{matrix} a_1, a_1 + m, a_3, a_4 \\ b_1, b_2, b_3 \end{matrix} ; z \right) \\ &= (-z)^{-a_1} \sum_{\nu=0}^{m-1} \frac{z^{-\nu}}{\nu!} \Gamma \left[\begin{matrix} m - \nu, a_3 - a_1 - \nu, a_4 - a_1 - \nu, a_1 + \nu \\ b_1 - a_1 - \nu, b_2 - a_1 - \nu, b_3 - a_1 - \nu \end{matrix} \right] z^{-\nu} \\ &+ (-z)^{-a_1} \sum_{\nu=m}^{\infty} \frac{(-1)^m (-z)^{-\nu}}{\nu! (\nu - m)!} \Gamma \left[\begin{matrix} a_1 + \nu, a_3 - a_1 - \nu, a_4 - a_1 - \nu \\ b_1 - a_1 - \nu, b_2 - a_1 - \nu, b_3 - a_1 - \nu \end{matrix} \right] [A + \log(-z)], \quad (2.1) \end{aligned}$$

where m is a non-negative integer, $a_1 \neq 0, -1, -2, \dots; |\arg(-z)| < \pi$;

$$\begin{aligned} A &= \psi(\nu + 1) + \psi(\nu + 1 - m) + \psi(a_3 - a_1 - \nu) + \psi(a_4 - a_1 - \nu) \\ &\quad - \psi(a_1 + \nu) - \psi(b_1 - a_1 - \nu) - \psi(b_2 - a_1 - \nu) - \psi(b_3 - a_1 - \nu); \quad (2.2) \end{aligned}$$

and $\psi(z) = d\{\log \Gamma(z)\}/dz$.

For $a_4 = b_3$ and $a_3 = b_2$, the equation (2.1) gives rise to a known result for the Gauss function ${}_2F_1$ [1, p.63].

3. Second Expansion

When three of the upper parameters differ by integers, there holds the formula:

$$\Gamma \left[\begin{matrix} a_1, a_1 + m, a_1 + m + n, a_2 \\ b_1, b_2, b_3 \end{matrix} \right] {}_4F_3 \left(\begin{matrix} a_1, a_1 + m, a_1 + m + n, a_2 \\ b_1, b_2, b_3 \end{matrix} ; z \right)$$

$$\begin{aligned}
&= (-z)^{-a_1} \sum_{\nu=0}^{m-1} \frac{z^{-\nu}}{\nu!} \Gamma \left[\begin{matrix} m-\nu, m+n-\nu, a_1+\nu, a_2-a_1-\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \\
&\quad + (-z)^{-a_1} (-1)^m \sum_{\nu=m}^{m+n-1} \frac{(-z)^{-\nu}}{\nu!(\nu-m)!} \Gamma \left[\begin{matrix} m+n-\nu, a_1+\nu, a_2-a_1-\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \\
&\quad \cdot \{\theta + \log(-z)\} \\
&\quad + (-z)^{-a_1} (-1)^n \sum_{\nu=m+n}^{\infty} \frac{z^{\nu}}{\nu!(\nu-m)!(\nu-m-n)!} \\
&\quad \cdot \Gamma \left[\begin{matrix} a_1+\nu, a_2-a_1-\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \left[B^2 + B' + 2 \log(-z) B + \{\log(-z)\}^2 \right], \quad (3.1)
\end{aligned}$$

where m and n are non-negative integers $a_1, a_1+m \neq 0, -1, -2, \dots$; and $|\arg(-z)| < \pi$;

$$\begin{aligned}
\theta &= \psi(\nu+1) + \psi(\nu-m+1) + \psi(m+n-\nu) - \psi(a_2-a_1-\nu) \\
&\quad - \psi(a_1+\nu) - \psi(b_1-a_1-\nu) - \psi(b_2-a_1-\nu) - \psi(b_3-a_1-\nu); \quad (3.2)
\end{aligned}$$

$$B = \theta + \psi(\nu-m-n+1) - \psi(m+n-\nu); \quad (3.3)$$

$$\begin{aligned}
B' &= 3\psi'(1) + 3\zeta(2,1) - \zeta(2,\nu+1) - \zeta(2,\nu-m+1) - \zeta(2,\nu-m-n+1) \\
&\quad - \psi'(a_2-a_1-\nu) + \psi'(a_1+\nu) \\
&\quad - \psi'(b_1-a_1-\nu) - \psi'(b_2-a_1-\nu) - \psi'(b_3-a_1-\nu). \quad (3.4)
\end{aligned}$$

Here $\zeta(\cdot, \cdot)$ is the generalized Zeta function defined by

$$\zeta(s, \nu) = \sum_{n=0}^{\infty} (\nu+n)^{-s}, \quad \nu \neq 0, -1, -2, \dots \quad (3.5)$$

For a detailed account of generalized Zeta function $\zeta(s, \nu)$, the reader is referred to the monograph by A. Erdélyi *et al.* [1].

4. Third Expansion

When four of the upper parameters differ by integers, the following result holds:

$$\begin{aligned}
&\Gamma \left[\begin{matrix} a_1, a_1+m, a_1+m+n, a_1+m+n+p \\ b_1, b_2, b_3 \end{matrix} \right] \\
&\quad \cdot {}_4F_3 \left(\begin{matrix} a_1, a_1+m, a_1+m+n, a_1+m+n+p \\ b_1, b_2, b_3 \end{matrix} ; z \right)
\end{aligned}$$

$$\begin{aligned}
&= (-z)^{-a_1} \sum_{\nu=0}^{m-1} \Gamma \left[\begin{matrix} m-\nu, m+n-\nu, m+n+p-\nu, a_1+\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] z^{-\nu} \\
&\quad + (-z)^{-a_1} \sum_{\nu=m}^{m+n-1} \frac{(-1)^m z^{-\nu}}{\nu!(\nu-m)!} \Gamma \left[\begin{matrix} m+n-\nu, m+n+p-\nu, a_1+\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \\
&\quad \cdot [C + \log(-z)] \\
&\quad + (-z)^{-a_1} \sum_{\nu=m+n}^{m+n+p-1} \frac{(-1)^n z^{-\nu}}{2!\nu!(\nu-m)!(\nu-m-n)!} \\
&\quad \cdot \Gamma \left[\begin{matrix} m+n+p-\nu, a_1+\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \\
&\quad \cdot [D' + D^2 + 2D \log(-z) + \{\log(-z)\}^2] \\
&\quad + (-z)^{-a_1} \sum_{\nu=m+n+p}^{\infty} \frac{(-1)^{m+p} (-z)^{-\nu}}{\nu!(\nu-m)!(\nu-m-n)!(\nu-m-n-p)!} \\
&\quad \cdot \Gamma \left[\begin{matrix} a_1+\nu \\ b_1-a_1-\nu, b_2-a_1-\nu, b_3-a_1-\nu \end{matrix} \right] \\
&\quad \cdot [(E^3 + 3E'E + E'') + 3\log(-z)(E' + E^2) + 3\{\log(-z)\}^2 E + \{\log(-z)\}^3], \quad (4.1)
\end{aligned}$$

where m, n and p are non-negative integers, $a_1, a_1+m, a_1+m+n \neq 0, -1, -2, \dots, |\arg(-z)| < \pi$,

$$\begin{aligned}
C &= \psi(\nu+1) + \psi(\nu-m+1) + \psi(m+n-\nu) + \psi(m+n+p-\nu) - \psi(a_1+\nu) \\
&\quad - \psi(b_1-a_1-\nu) - \psi(b_2-a_1-\nu) - \psi(b_3-a_1-\nu); \quad (4.2)
\end{aligned}$$

$$D = C + \psi(\nu-m-n+1) - \psi(m+n-\nu); \quad (4.3)$$

$$\begin{aligned}
D' &= 3\psi'(1) + 3\zeta(2,1) - \zeta(2,\nu+1) - \zeta(2,\nu-m+1) - \zeta(2,\nu-m-n+1) \\
&\quad - \psi'(m+n+p-\nu) + \psi'(a_1+\nu) \\
&\quad - \psi'(b_1-a_1-\nu) - \psi'(b_2-a_1-\nu) - \psi'(b_3-a_1-\nu); \quad (4.4)
\end{aligned}$$

$$E = D + \psi(\nu-m-n-p+1) - \psi(m+n+p-\nu); \quad (4.5)$$

$$E' = D' + \psi'(1) + \zeta(2,1) + \psi'(m+n+p-\nu) - \zeta(2,\nu-m-n-p+1); \quad (4.6)$$

$$\begin{aligned}
E'' &= 2[2\psi''(1) + 4\zeta(3,1) - \zeta(3,\nu+1) - \zeta(3,\nu-m+1) \\
&\quad - \zeta(3,\nu-m-n+1) - \zeta(3,\nu-m-n-p+1)] \\
&\quad - \psi''(a_1+\nu) - \psi''(b_1-a_1-\nu) - \psi''(b_2-a_1-\nu) - \psi''(b_3-a_1-\nu). \quad (4.7)
\end{aligned}$$

Remark 1. It may be mentioned here that the results given earlier by Saxena and Kalla [26, pp.41-43] follow from (2.1) and (3.1).

Remark 2. Though the formulae (2.1), (3.1) and (4.1) are special cases of a general one for ${}_pF_q$ listed in [7, §7.2.3.78], they are still sufficiently general and unify known results in [1, p.63], [27, pp.41-43] etc.

Acknowledgement. The present investigation was supported, in part, by the Ministry of Education of Japan under Grant 09640238 and the University Grants Commission of India. Thanks are due to Professor H.M. Srivastava for giving some useful suggestions.

References

- [1] A. Erdélyi, W. Magnus, F. Oberhettiger and F.G. Tricomi: *Higher Transcendental Functions*, Vol.1, McGraw-Hill, New York, 1953.
- [2] W. Magnus, F. Oberhettiger and R.P. Soni: *Formulas and Theorems for the Special Functions of Mathematical Physics*, Third Enlarged Ed., Springer-Verlag, Berlin-Heidelberg, 1966.
- [3] A.M. Mathai: A few remarks on the exact distributions of certain multivariate statistics - II. *Multivariate Statistical Inference*, pp. 169-181, Kabe, D.G. and Gupta, R.P. (Eds.), North Holland, Amsterdam, 1972.
- [4] A.M. Mathai and H.J. Haubold: *Modern Problems in Nuclear and Neutrino Astrophysics*, Akademie-Verlag, Berlin, 1988.
- [5] A.M. Mathai and R.K. Saxena: *Generalized Hypergeometric Function with Applications in Statistics and Physical Sciences*, Lecture Notes Math., Vol. 348, Berlin, 1973.
- [6] A.M. Mathai and R.K. Saxena: *The H-function with Applications in Statistics and Other Disciplines*, Halsted Press, New York-London-Sydney-Toronto, 1978.
- [7] A.P. Prudnikov, Yu.A. Brychkov and O.I. Marichev: *Integral and Series*, Vol.3, *More Special Functions*, Gordon and Breach, New York-London, 1989.
- [8] E.D. Rainville: *Special Functions*, Chelsea Pub., New York, 1960.
- [9] M. Saigo: On a property of the Appell hypergeometric function F_1 , *Math. Rep. Kyushu Univ.* **12**(1980), 63-67.
- [10] M. Saigo: On properties of the Appell hypergeometric functions F_2 and F_3 and the generalized Gauss function ${}_3F_2$, *Bull. Centr. Res. Inst., Fukuoka Univ.* **66**(1983), 27-32.
- [11] M. Saigo: On properties of the Lauricella hypergeometric function F_D , *Bull. Centr. Res. Inst., Fukuoka Univ.* **104**(1988), 13-31.
- [12] M. Saigo: On properties of hypergeometric functions of three variables, F_M and F_G , *Rend. Circ. Mat. Palermo, Serie II* **37**(1988), 449-468.
- [13] M. Saigo: On behaviors of hypergeometric series of one, two, three and n -variables near boundaries of their convergence regions of logarithmic case, (Kyoto, 1989), *Sûrikaiseiki-kenkyûsho Kôkyûroku* **714**(1990), 91-109.
- [14] M. Saigo: The asymptotic behaviors of triple hypergeometric series F_G, F_K, F_N and F_R near boundaries of their convergence regions (Kyoto, 1989), *Sûrikaiseiki-kenkyûsho Kôkyûroku* **714**(1990), 110-126.

- [15] M. Saigo: Asymptotic behaviors of the Horn hypergeometric series $G_1, G_2, H_1, H_2, H_4, H_6, H_7$ near boundaries of their convergence regions, *Integral Transform. Spec. Funct.* **4**(1996), 249-262.
- [16] M. Saigo, O.I. Marichev and Ngyen Thanh Hai: Asymptotic representations of Gaussian series ${}_2F_1$, Clausenian series ${}_3F_2$ and Appell series F_2 and F_3 near boundaries of their convergence regions, *Fukuoka Univ. Sci. Rep.* **19**(1989), 83-90.
- [17] M. Saigo and Ngyen Thanh Hai: The asymptotic representations of Appell series F_1 and Horn series H_3 near boundaries of their convergence regions, *Fukuoka Univ. Sci. Rep.* **21**(1991), 17-23.
- [18] M. Saigo and Ngyen Thanh Hai: Asymptotic representations of the Horn series G_1 and G_2 near boundaries of their convergence regions, *Fukuoka Univ. Sci. Rep.* **22**(1992), 1-9.
- [19] M. Saigo and H.M. Srivastava: The behaviors of the Appell double hypergeometric series F_4 and certain Lauricella triple hypergeometric series near the boundaries of their convergence regions, *Fukuoka Univ. Sci. Rep.* **19**(1989), 1-10.
- [20] M. Saigo and H.M. Srivastava: The asymptotic behaviour of Lauricella's second hypergeometric series $F_B^{(n)}$ in n variables near the boundary of its convergence region, *Fukuoka Univ. Sci. Rep.* **19**(1989), 71-82.
- [21] M. Saigo and H.M. Srivastava: The behaviors of the zero-balanced hypergeometric series ${}_pF_{p-1}$ near the boundary of its convergence region, *Proc. Amer. Math. Soc.* **110**(1990), 71-76.
- [22] M. Saigo and H.M. Srivastava: Some asymptotic formulas exhibiting the behaviors of the triple hypergeometric series F_S and F_T near the boundaries of their convergence regions, *Funk. Ekvac.* **34**(1991), 423-448.
- [23] M. Saigo and H.M. Srivastava: A theorem on the asymptotic behavior of Lauricella's first hypergeometric series $F_A^{(n)}$ near the boundary of its convergence region, *Complex Analysis and Generalized Functions*, Varna, September 15-21 (1991), 258-267.
- [24] M. Saigo and H.M. Srivastava: The asymptotic behaviour of the Srivastava hypergeometric series H_C near the boundary of its convergence region, *Kyungpook Math. J.* **32**(1992), 153-158.
- [25] M. Saigo and H.M. Srivastava: The behaviour of the fourth type of Lauricella's hypergeometric series in n variables near the boundaries of its convergence region, *J. Austral. Math. Soc., Ser. A* **57**(1994), 281-294.
- [26] R.K. Saxena and S.L. Kalla: Expansion of a ${}_3F_2$ when the upper parameters differ by an integer, *Rev. Tec. Ing. Univ. Zulia* **9**(1986), 41-43.